

Seismology of triple-mode classical Cepheids of the Large Magellanic Cloud

P. Moskalik¹ and W. A. Dziembowski^{2,1}

¹ Copernicus Astronomical Center, ul. Bartycka 18, 00-716 Warsaw, Poland

² Warsaw University Observatory, Al. Ujazdowskie 4, 00-478 Warsaw, Poland

Received 5 October 2004/ Accepted 23 December 2004

Abstract. We interpret the three periods detected in OGLE LMC Cepheids SC3-360128 and SC5-338399 as corresponding to the first three overtones of radial pulsations. This interpretation imposes stringent constraints on parameters of the stars and on their evolutionary status, which could only be the first crossing of the instability strip. Evolutionary models reproducing measured periods exist only in a restricted range of metallicities ($Z = 0.004 - 0.007$). The models impose an upper limit on the extent of overshooting from the convective core. Absolute magnitude of each star is confined to a narrow interval. This allows to derive a new estimate of the distance to the LMC. We obtain $m - M$ ranging from $18^m 34$ to $18^m 53$, with a systematic difference between the two stars of about $0^m 13$. The rates of period change predicted by the models are formally in conflict with the derived observational limits, though the uncertainties of measured $\dot{P}$ may be underestimated. If the discrepancy is confirmed, it would constitute a significant challenge to the stellar evolution theory.

Key words. Stars: variables: Cepheids – Stars: oscillations – Stars: evolution – Galaxies: individual: LMC

1. Introduction

Triple-mode radial pulsators are extremely rare. For many years AC Andromedae, with the fundamental and the first two overtones simultaneously excited (Fitch & Szeidl 1976), was the only known object of this type. Recently, two new triple-mode variables have been identified among short period Cepheids of the Large Magellanic Cloud (Moskalik, Kołaczowski & Mizerski 2004, Moskalik & Kołaczowski 2005). Both stars, LMC SC3-360128 (hereafter Star 1) and LMC SC5-338399 (hereafter Star 2), have previously been known to pulsate in two radial modes, namely in the first and second overtones (Soszyński et al. 2000). The analysis of Moskalik & Kołaczowski (2005) clearly shows presence of the third independent periodicity. The period ratio of $P_3/P_2 = 0.840$, the same in both Cepheids, points towards identification of the newly detected mode as a third radial overtone. The two stars are the first known pulsators with three consecutive radial overtones simultaneously excited.

Even if certain forms of pulsation are rare, it does not mean that they are unimportant. The best example is the role of double-mode Cepheid data in stimulating the revision of stellar opacities. Each measured period of an identified mode yields an accurate constraint on stellar pa-

rameters. For our objects we have three such constraints. Another advantage is the membership of both Cepheids in the LMC. The stars have been and will be systematically monitored. Moreover, we have reasonably good estimate of their absolute luminosities. This estimate is helpful but not precise, because the distance to the LMC is still debated. In fact, we will show that seismological analysis of the two triple-mode Cepheids yields an independent distance determination.

Our aim is to extract maximum information on the two stars from all the available data. After summarizing observational information for the two objects (Section 2), we focus on the inference from the values of measured periods. Our analysis in Section 3 is similar to that of Kovács & Buchler (1994) for AC And. We conclude, that both triple-mode Cepheids must be in the evolutionary phase of the Hertzsprung gap crossing and we confront seismologically derived stellar parameters with predictions of the appropriate evolutionary models (Section 4). This further constrains the ranges of admissible stellar parameters. Section 5 is devoted to the LMC distance estimation implied by our models. In Section 6 we compare theoretically predicted rates of period change with the limits inferred from observations. Finally, in Section 7 we summarize our conclusions and discuss future observations.

Send offprint requests to: P. Moskalik,
e-mail: pam@camk.edu.pl

Table 1. Triple-mode Cepheids in the LMC

OGLE No.	P_1 [day]	P_2 [day]	P_3 [day]	V [mag]	I [mag]	$E(B - V)$ [mag]	$(V - I)_0$ [mag]	W_I [mag]
LMC SC3-360128	0.541279	0.436049	0.366301	17.53	16.98	0.134	0.38	16.13
LMC SC5-338399	0.579510	0.466624	0.392116	17.37	16.87	0.133	0.33	16.09

2. Data

Main observational properties of both triple-mode Cepheids are summarized in Table 1. Pulsation periods are determined from OGLE I -band photometry reduced with the DIA algorithm (Žebruň et al. 2001). All periods are accurate to better than 7×10^{-6} day. Their errors are not quoted here; we believe they are smaller than uncertainties of the model periods and, therefore, they are inconsequential for our analysis. The mean V -band and I -band magnitudes are taken from Soszyński et al. (2000). The average $E(B - V)$ reddening for the field of each Cepheid (Udalski et al. 1999) is used to obtain dereddened $(V - I)_0$ colours. Standard extinction coefficients are adopted: $A_V = 3.24 E(B - V)$, $A_I = 1.96 E(B - V)$. In the last column of the table we give Wessenheit index, $W_I = I - 1.55 \times (V - I)$, which is an extinction insensitive brightness indicator (Madore & Freedman 1991).

3. Constraints on stellar parameters from pulsation periods

The basic assumption behind our analysis is that the three periods detected in the triple-mode Cepheids correspond to the first three consecutive overtones of radial pulsations. The constraints on the stellar parameters are derived by equating the three measured periods $P_{n,o}$ with periods calculated in plausible stellar models, $P_{n,c}$. In this section we use only envelope models, extending from the photosphere down to the temperature of $\log T = 7.2$. This is the standard approach in studies of radial pulsations in giants. We checked that the calculated periods are sufficiently insensitive to the location of the inner boundary of the envelope. The model periods depend on six parameters. These are: the mass, M , luminosity, L , effective temperature, T_{eff} , the fractional abundances of hydrogen, X , and of elements heavier than helium, Z , and the mixing length parameter, α_{con} . The pulsation data yield for each star three equations of the form

$$P_{n,c}(M, L, T_{\text{eff}}, X, Z, \alpha_{\text{con}}) = P_{n,o}$$

and this is not enough to determine the values of all the unknown parameters. In principle, we could use mean magnitudes and colours of the stars, but these are uncertain. Therefore, we decided to find solutions for M , L and T_{eff} for selected X , Z , and α_{con} in admissible ranges, relying only on the measured periods and requiring that all three modes are linearly unstable at specified parameters. Comparison with the observed mean colours and magnitudes is done *a posteriori*.

Our static stellar models, both envelope models used here and the evolutionary models used in Section 4.2, are calculated with a modernized version of the Paczyński (1970) codes. The main improvements consist in the implementation of the OPAL equation-of-state and opacity data (Rogers, Swenson & Iglesias 1996; Iglesias & Rogers 1996). Pulsation periods are calculated with a modified version of the Dziembowski (1977) code.

There are uncertainties in the calculated periods. We rely on the linear nonadiabatic approximation. We believe this is the most severe simplification. Comparison of periods derived from linear and from nonlinear codes has been published only for monoperiodic RR Lyr models (e.g. Kovács & Buchler 1988) and long period Cepheid models (e.g. Moskalik & Buchler 1991; Fokin 1994). It is not clear how to scale these results to higher overtones in multiperiodic $\sim 3M_{\odot}$ Cepheids.

Errors may result from using the mixing length theory to calculate convective flux in our static models and ignoring its Lagrangian perturbation in pulsating models. We believe that this is less essential and that our calculations made with $\alpha_{\text{con}} = 0$ and $\alpha_{\text{con}} = 1$ reflect well the uncertainty. The use of the Eddington approximation in the optically shallow layers is also an inadequacy but of secondary importance for the present application.

Simultaneous fit to three radial mode periods can be achieved for several different sets of parameters (*e.g.* AC And; Kovács & Buchler 1994). In order to assure that we find *all* the solutions we apply the following procedure:

1. first, we select the chemical composition (X , Z) and the mixing length α_{con} . Keeping these three parameters fixed, we calculate a grid of stellar models covering the mass range of $2 < M/M_{\odot} < 5$ and luminosity range of $2 < \log(L/L_{\odot}) < 3$.
2. at every point of the (M, L) grid we find a model which fits the observed period of the first overtone, $P_{1,o}$. This can be easily achieved by adjusting the last remaining parameter of the model, namely its effective temperature. The solution for T_{eff} is unique. The model fitting P_1 does not necessarily reproduce the other two periods. We calculate period mismatches $\Delta P_2 = P_{2,c} - P_{2,o}$ and $\Delta P_3 = P_{3,c} - P_{3,o}$.
3. we find on the (M, L) plane the lines of $\Delta P_2 = 0$. Models located on these lines reproduce *both* P_1 and P_2 . Similarly, we find the lines of $\Delta P_3 = 0$. The values of (M, L) at which the two types of lines cross each other define the models which simultaneously reproduce *all three periods*.




Fig. 1. Simultaneous fit to three pulsation periods observed in Star 2. Three panels display models calculated for $\alpha_{\text{con}} = 0$, $X = 0.70$ and metallicities of $Z = 0.009$, $Z = 0.007$ and $Z = 0.005$. Filled and open circles denote lines of $\Delta P_2 = 0$ and $\Delta P_3 = 0$, respectively. Different triple-mode solutions are marked A, B and C. Dotted area is the domain where all three modes are linearly unstable.

Table 2. Seismic envelope models for Star 1

α_{con}	X	Z	$M/M_{\odot}$	$\log(L/L_{\odot})$	$\log T_{\text{eff}}$ [K]	$(V - I)$ [mag]	$m - M$ [mag]
0	0.74	0.006	3.29	2.311	3.814	0.50	18.43
			3.63	2.403	3.829	0.43	18.52
0	0.74	0.005	3.15	2.358	3.828	0.44	18.42
			(3.67)	(2.487)	(3.849)		
1	0.74	0.007	3.48	2.299	3.807	0.52	18.47
			3.67	2.359	3.818	0.48	18.52
1	0.74	0.006	3.21	2.316	3.817	0.48	18.42
			3.68	2.423	3.834	0.41	18.53
1	0.74	0.005	3.14	2.376	3.833	0.42	18.41
			3.59	2.461	3.845	0.37	18.51
1	0.74	0.004	3.14	2.433	3.847	0.36	18.42
			(3.60)	(2.547)	(3.865)		
0	0.70	0.008	3.28	2.195	3.787	0.60	18.39
			3.48	2.263	3.800	0.55	18.45
0	0.70	0.007	3.10	2.215	3.795	0.57	18.37
			3.63	2.366	3.822	0.46	18.50
0	0.70	0.006	2.98	2.279	3.813	0.50	18.36
			3.66	2.437	3.839	0.39	18.52
0	0.70	0.005	2.88	2.346	3.832	0.42	18.35
			(3.66)	(2.506)	(3.856)		
1	0.70	0.007	3.21	2.240	3.799	0.56	18.40
			3.67	2.388	3.827	0.44	18.51
1	0.70	0.006	2.96	2.300	3.819	0.47	18.36
			3.65	2.440	3.840	0.39	18.51
1	0.70	0.005	2.90	2.362	3.836	0.40	18.35
			(3.61)	(2.487)	(3.852)		
1	0.70	0.004	2.86	2.410	3.848	0.35	18.35
			(3.63)	(2.575)	(3.873)		

Table 3. Seismic envelope models for Star 2

α_{con}	X	Z	$M/M_{\odot}$	$\log(L/L_{\odot})$	$\log T_{\text{eff}}$ [K]	$(V - I)$ [mag]	$m - M$ [mag]
0	0.70	0.008	3.45	2.333	3.807	0.52	18.52
			3.65	2.382	3.815	0.49	18.57
0	0.70	0.007	3.29	2.367	3.818	0.48	18.50
			3.74	2.461	3.833	0.42	18.60
0	0.70	0.006	3.21	2.424	3.834	0.41	18.49
			(3.74)	2.525	3.848)		
1	0.70	0.008	3.41	2.337	3.809	0.51	18.51
			3.74	2.416	3.823	0.46	18.58
1	0.70	0.007	3.27	2.385	3.824	0.45	18.49
			3.75	2.469	3.836	0.40	18.59
1	0.70	0.006	3.25	2.438	3.837	0.40	18.50
			3.67	2.504	3.845	0.36	18.59
1	0.70	0.005	3.26	2.489	3.849	0.35	18.51
			(3.64)	2.563	3.860)		
0	0.72	0.007	3.52	2.397	3.822	0.46	18.54
			3.66	2.430	3.827	0.44	18.58
0	0.72	0.006	3.38	2.438	3.834	0.41	18.53
			(3.72)	2.508	3.845)		
1	0.72	0.007	3.42	2.399	3.824	0.45	18.53
			3.73	2.456	3.832	0.42	18.60
1	0.72	0.006	3.45	2.458	3.838	0.39	18.54
			3.59	2.482	3.841	0.38	18.57

In Fig. 1 we display the results of our procedure applied to Star 2, for $\alpha_{\text{con}} = 0$, $X = 0.70$ and three different metallicity values of $Z = 0.009, 0.007$ and 0.005 . The filled and open circles define the lines of $\Delta P_2 = 0$ and $\Delta P_3 = 0$, respectively. The dotted area in each plot corresponds to models with all three modes linearly unstable. We recall here, that in the absence of resonances (as in our case), the linear excitation of all modes is the necessary condition for the triple-mode pulsations to exist. Among all models matching the observed periods, only those located in the dotted area correspond to acceptable solutions.

As we see from Fig. 1, the problem of matching periods of a triple-mode Cepheid can have up to three different solutions. We call them *A*, *B* and *C*. Solution *A* exists for all choices of X , Z and α_{con} , but all three modes are always linearly stable. We will not discuss this solution any further. Solutions *B* and *C* can exist only for metallicities below a certain critical value. This is clearly seen in Fig. 1: for $Z = 0.009$ the lines $\Delta P_2 = 0$ and $\Delta P_3 = 0$ are close to each other, but do not cross. As the value of Z is lowered, the two lines come closer and finally cross each other, generating solutions *B* and *C*. For the case displayed in Fig. 1, this happens for $Z \sim 0.0082$. When Z is lowered further, first solution *C* and then solution *B* migrates out of the dotted area. This sets a *lower limit* for the metallicity. Below this limit solutions *B* and *C* do exist, but do not correspond to triple-mode pulsations, because at least

one of the modes is linearly stable. For the case of Fig. 1 this lower limit is at $Z \sim 0.0056$.

We have performed search for solutions for both triple-mode Cepheids, exploring the range of metallicities from $Z = 0.004$ to 0.010 (in some cases to 0.020) and of hydrogen abundances from $X = 0.70$ to 0.74 . For each chemical composition we have calculated models with $\alpha_{\text{con}} = 0$ and with $\alpha_{\text{con}} = 1$. The results of this survey are summarized in Table 2 for Star 1 and in Table 3 for Star 2. We list only those combinations of X , Z and α_{con} for which at least one acceptable solution exists. For low metallicities, solution *C* may already fall outside the "all-modes-unstable" domain, in such a case it is given in parenthesis. For all acceptable solutions we provide the values of $(V - I)$ colour and the distance modulus to the star, $m - M$. The latter is obtained by comparing Wessenheit index, W_I , derived from the models with the mean observed values, as given in Table 1. The conversion from model parameters to V -band and I -band magnitudes is done with tabular data based on Kurucz's (1998) stellar atmosphere models.

The triple-mode pulsations strongly constrain parameters of the studied Cepheids. We find, that metallicities of both stars have to be in the range of $Z = 0.0035 - 0.008$, with limiting values somewhat dependent on X and assumed α_{con} . For Star 1, fitting the observed periods is possible for all values of X , but for Star 2 hydrogen abundance of $X \leq 0.72$ is required. The inferred stellar masses

are limited to ranges of $2.8 < M/M_\odot < 3.7$ for Star 1 and $3.2 < M/M_\odot < 3.8$ for Star 2. The respective distance moduli are in the ranges of $18^m35 - 18^m53$ and $18^m49 - 18^m60$.

In the next section we will show that the constraints imposed by triple-mode pulsations can be considerably tightened by combining the seismic envelope models with the appropriate evolutionary tracks.

4. Evolutionary models

4.1. Evolutionary status of the stars

The very short periods and low luminosities of the triple-mode Cepheids cannot be explained with the current models of stars in the phase of central helium burning. Whether or not a star enters the instability strip during this evolutionary phase depends on the extent of the *blue loop*. The latter is mainly determined by the mass and the metallicity of the star. Alibert et al. (1999) find that at $Z = 0.004$ their models with masses of $3.0 - 3.5M_\odot$ do enter the instability strip. However, the minimum value of P_1 , which occurs for $M = 3M_\odot$ is about 0^d9 . At $Z = 0.01$ they find models with blue loop extending to the instability strip only at $M \geq 4M_\odot$. Such models have, naturally, much higher luminosities and pulsation periods than our stars. Alibert et al. (1999) assumed negligible overshooting in their calculations.

The effect of finite overshooting was taken into account in an extensive set of models of Girardi et al. (2000). In this case, the parameters we have derived for the triple-mode Cepheids are even more difficult to reconcile with the models in the helium burning phase. At $Z = 0.004$ and $M = 3M_\odot$ the loop does not extend to the instability strip. At $M = 3.5M_\odot$ the loop ends within the strip, but then the minimum value of P_1 is about 1^d15 . Again, increasing Z only worsens the discrepancy.

Determination of the extent of the blue loop is a subtle problem. Here we find the largest discrepancies between various authors. A more robust result is the difference of $\Delta \log L \sim 0.2$ between the first and the second crossing of the instability strip by stars in the $3 - 4M_\odot$ mass range. The $\log L$ values we find for the triple-mode Cepheids indicate that these two objects are caught during the short phase of the first crossing of the strip.

4.2. Models in the first crossing of the instability strip

We have computed a number of evolutionary tracks covering the Hertzsprung gap crossing phase. From each track, we may determine L in the moment when the first overtone period is equal to $P_{1,o}$. The value of L is virtually independent of α_{con} , but depends strongly on α_{ov} , which is the ratio of the overshooting distance from the convective core to the pressure distance scale calculated at the edge of the core. From our model calculations we find




Fig. 2. Evolutionary $M - L$ relation for models with $P_1 = 0^d541$. Lines are derived for stellar models undergoing first crossing of the instability strip. Models are calculated assuming no overshooting from the convective core ($\alpha_{\text{ov}} = 0$) during the main sequence evolution. Symbols represent seismic solutions for Star 1, derived with (non-evolutionary) envelope models (see Table 2). Filled and open symbols correspond to $\alpha_{\text{con}} = 1$ and 0 , respectively. Pentagons correspond to $X = 0.70, Z = 0.006$, circles to $X = 0.74, Z = 0.006$, triangles to $X = 0.74, Z = 0.005$, and square to $X = 0.74, Z = 0.007$ (there is no solution for $\alpha_{\text{con}} = 0$ in the last case).

$$\left(\frac{\partial \log L}{\partial \alpha_{\text{ov}}} \right)_{M, X, Z, P_1} \approx 0.6$$

At specified values of M, X, Z and P_1 , the minimum value of L corresponds to $\alpha_{\text{ov}} = 0$

In Fig. 2 we display the $M - L$ relations for no overshooting evolutionary models of Star 1 ($P_1 = 0^d541$). Different lines correspond to various chemical compositions. In the same plot we also display parameters of several seismic models of Star 1, also for different values of X and Z (see Table 2). We plot only models corresponding to solution *B*, because only for this solution agreement between seismic and evolutionary models can be reached. The lower panel of the plot explores the effect of varying stellar metallicity. For $Z = 0.005$ the seismic models do not agree with $\alpha_{\text{ov}} = 0$ evolutionary tracks. The agreement may be easily reached if overshooting of $\alpha_{\text{ov}} \sim 0.1$ is allowed, which shifts the evolutionary $M - L$ relation up. For $Z = 0.006$ the seismic models and the evolutionary tracks are almost in agreement. For $Z = 0.007$ the seismic model is placed below the corresponding evolutionary $M - L$ line and no agreement between the two is possible. Fig. 2 illustrates the general pattern of model behaviour: with X, α_{con} and α_{ov} fixed, the exact match between seis-

Table 4. Triple-mode Cepheid models consistent with no overshooting evolutionary tracks

X	α_{con}	Z	$M/M_{\odot}$	$\log(L/L_{\odot})$	$\log T_{\text{eff}}$ [K]	$(V-I)$ [mag]	$m-M$ [mag]	$\dot{P}/P$ [Myr $^{-1}$]
Star 1								
0.70	0	0.0061	2.99	2.273	3.811	0.51	18.36	1.51
	1	0.0063	3.02	2.280	3.813	0.50	18.37	1.53
0.74	0	0.0057	3.25	2.323	3.818	0.48	18.43	1.65
	1	0.0062	3.26	2.308	3.814	0.50	18.43	1.61
Star 2								
0.70	0	0.0065	3.24	2.394	3.826	0.44	18.49	2.15
	1	0.0069	3.26	2.392	3.826	0.44	18.49	2.18
0.72	0	0.0061	3.39	2.432	3.832	0.42	18.53	2.11
	1	—	—	—	—	—	—	—

mic models and evolutionary tracks can be reached only at one specific metallicity value. The matching values of Z becomes lower, when the overshooting distance α_{ov} is increased.

The upper panel of Fig. 2 shows that match between seismic and evolutionary models is possible for any reasonable value of X , but yielding different values of stellar mass: $M \sim 3.2M_{\odot}$ for $X = 0.74$ and $M \sim 3.0M_{\odot}$ for $X = 0.70$. Interestingly, metallicities of matching models are almost the same independently of X .

In Table 4 we list parameters of the two stars resulting from matching seismic models to the evolutionary tracks calculated for $\alpha_{\text{ov}} = 0$. The values obtained with two choices of α_{con} give some assessment of uncertainty due to poor knowledge of convection. We regard the parameters derived at higher X to be more likely, because we do not expect any significant He enrichment for LMC stars with $M \sim 3M_{\odot}$. In the last two columns we provide the values of $(V-I)$ colour and of the distance modulus to the stars, obtained in same way as in Tables 2 and 3. Bearing in mind uncertainty in the mean dereddened colour, which according to Udalski (private communication) is about 0^m1, we regard the agreement in $(V-I)$ to be quite satisfactory. The agreement may be even better, if smaller LMC redenings of Subramaniam (2004) are used. We will discuss the inferred values of the distance modulus later in this paper. In the last column of Table 4 we list the calculated rates of period change. These numbers will be discussed later, too.

We stress, that with exception of $\dot{P}/P$, all the values listed in Table 4 are derived from the envelope models. The requirement of matching the seismic envelope models and the evolutionary tracks was used only to a) reject solution C and b) to find the value of Z at which the exact match is achieved.

4.3. Constraints on the extent of overshooting

At fixed X and α_{con} , there is a minimum metallicity value allowed by seismic models. This lower limit for Z

corresponds to an *upper limit* for the overshooting distance, α_{ov} . In Table 5 we give the maximum overshooting distances for both triple-mode Cepheids. We obtain a stronger constraint for Star 2, especially if the higher value of X is adopted. Despite very similar masses, we cannot argue that the two stars should have similar α_{ov} . They may well differ in the rotation rate and rotation induced mixing can mimic the convective overshooting. In any case, the limit of $\alpha_{\text{ov}} \leq 0.12$ obtained for Star 2 is very interesting.

The triple-mode Cepheids can be quite important for constraining element mixing in moderate mass stars. To demonstrate this point, we list in Table 5 also the stellar parameters of the two Cepheids derived at the extreme values of α_{ov} . These can be compared with parameters derived for $\alpha_{\text{ov}} = 0$ (Table 4). As overshooting is increased, the matching models become less metal abundant and bluer. Thus, the limits on the extent of overshooting can be further improved once we precisely measure either metallicities or the mean dereddened colours of the two stars. We assume here, that the composition in the envelope and in the photosphere is the same, which is valid unless the star is a fast rotator.

5. The distance to LMC

The values of distance moduli listed in Tables 4 and 5 are all between 18^m34 and 18^m53, *i.e.* well within currently considered range for the LMC. They are indeed very close to estimates based on analysis of RRd stars, which yields 18^m53 according to Kovács (2000b) and 18^m42 according to Popielski et al. (2000). From the double-mode Cepheid data, Kovács (2000a) obtained the value of 18^m54. All these determinations rely on fitting the observed pulsation periods, but for different objects and in different ways.

For the same model parameters (X , α_{con}), there is a systematic difference of about 0^m13 between distance moduli derived for Star 1 and Star 2. We believe, that large part of this difference can be attributed to W_I measurement errors. Indeed, judging from faint Cepheids

Table 5. Triple-mode Cepheid models for maximum overshooting allowed by seismic constraints

X	α_{con}	Z	α_{ov}	$M/M_{\odot}$	$\log(L/L_{\odot})$	$\log T_{\text{eff}}$ [K]	$(V - I)$ [mag]	$m - M$ [mag]	$\dot{P}/P$ [Myr $^{-1}$]
Star 1									
0.70	0	0.0043	0.33	2.81	2.393	3.845	0.36	18.34	1.69
	1	0.0036	0.33	2.84	2.428	3.853	0.33	18.34	1.81
0.74	0	0.0045	0.22	3.07	2.392	3.838	0.40	18.40	1.61
	1	0.0038	0.19	3.14	2.443	3.849	0.35	18.42	1.75
Star 2									
0.70	0	0.0056	0.12	3.18	2.450	3.841	0.38	18.49	2.07
	1	0.0049	0.07	3.26	2.493	3.850	0.34	18.51	2.26
0.72	0	0.0058	0.05	3.35	2.452	3.838	0.39	18.52	2.12
	1	—	—	—	—	—	—	—	—

($W_I > 15^{\text{m}}0$) with independent photometry from two OGLE fields, the error of the mean Wesenheit index is about $0^{\text{m}}06$ *on average*, but sometimes can exceed $0^{\text{m}}09$. We cannot exclude, however, that some part of the $0^{\text{m}}13$ difference might be real. As shown by Subramaniam (2003), the $m - M$ displays considerable variation even within the LMC bar.

We want to stress, that distances of the triple-mode Cepheids can be estimated very accurately once photometric precision is improved. As seen from Tables 4 and 5, the total range of $m - M$ allowed by seismic models is only $0^{\text{m}}09$ for Star 1 and $0^{\text{m}}04$ for Star 2. In principle, the individual distances of the two stars can be determined with, respectively, $\pm 2\%$ and $\pm 1\%$ accuracy.

6. Period changes

The first crossing of the instability strip is by two orders of magnitude faster than either the second or the third crossing. Measuring the rates of period change should, therefore, provide a crucial test for the triple-mode Cepheid models.

The rates of period change, $\dot{P}/P$, inferred from our models are given in the last column of Tables 4 and 5. They are about 2 Myr^{-1} , somewhat below this value for Star 1 and somewhat above it for Star 2. The rates are calculated for the first overtone, but they are approximately the same for all modes. With such values of $\dot{P}/P$, the $(O - C)$ should reach 6 – 10% of the pulsation period in ten years.

We do not have yet ten years worth of data at our disposal, but we will try to measure $\dot{P}$ with photometry already at hand. Our method consists in the least square fitting of the photometric measurements with the Fourier sum of the form

$$I(t) = \langle I \rangle + \sum_{i=1}^3 A_i \cos\left[\left(\omega_i + \frac{1}{2}\dot{\omega}_i t\right)t + \phi_i\right] + \\ + \text{harmonics} + \text{combination terms}$$

where $\omega_i = 2\pi/P_i$ are pulsation frequencies and $\dot{\omega}_i = -2\pi\dot{P}_i/P_i^2$ are the rates of frequency change (see *e.g.* Kepler 1993 for derivation). The nonlinear least square fit is performed with the Marquardt algorithm (Press et al. 1986). In the following, we will discuss only the rates of period change for the first overtone. The other two modes have much lower amplitudes and consequently their $\dot{P}$ is determined with 3-10 times larger errors.

The method, when applied to DIA-reduced OGLE I -band photometry yields $\dot{P}_1/P_1 = +1.7 \pm 3.7 \text{ Myr}^{-1}$ for Star 1 and $\dot{P}_1/P_1 = +5.3 \pm 6.4 \text{ Myr}^{-1}$ for Star 2. Clearly, the OGLE timebase of ~ 1200 day is not sufficient to obtain significant $\dot{P}$ measurement.

A considerably longer timebase (~ 2700 day) is offered by MACHO photometry (Allsman & Axelrod 2001). Both triple-modes Cepheids have been observed by MACHO Project (LMC SC3-360128=MACHO 77.8029.310; LMC SC5-338399=MACHO 80.7074.120) and over 1500 measurements have been accumulated for each of them. Unfortunately, MACHO data are of much lower quality. We have decided to perform initial filtering of MACHO photometry, by rejecting a) all outlying measurements deviating by more than 5σ from average brightness of the star and b) all measurements with photometric errors (given by MACHO) larger than 5 times the average for the dataset. Further data clipping is done during the fitting procedure, when all points deviating by more than 5σ from the fit are also rejected. For the final determination of $\dot{P}$ we use the MACHO B -band photometry. The analysis yields

$$\dot{P}_1/P_1 = -1.66 \pm 0.83 \text{ Myr}^{-1} \quad \text{for Star 1}$$

$$\dot{P}_1/P_1 = -2.28 \pm 0.94 \text{ Myr}^{-1} \quad \text{for Star 2}$$

For both Cepheids $\dot{P}/P$ is determined only with 2σ accuracy. Therefore, we do not consider the above values to be significant detections, but rather to provide observational limits on $\dot{P}$. The results, nevertheless, are very interesting, because these limits seem to be in conflict with the

model predictions. The measured $\dot{P}/P$ values differ from theoretical ones (Tables 4 and 5) by 3.8σ for Star 1 and by 4.6σ for Star 2, respectively. If real, such a discrepancy would mean serious troubles for the evolutionary models. However, we should treat this result with a grain of salt. The quoted errors of $\dot{P}/P$ are formal statistical errors resulting from the nonlinear least square procedure. Such errors very often underestimate the true uncertainties of the fitted parameters. This is especially true in the case of multiperiodic pulsations (Costa, Kepler & Winget 1999; Costa & Kepler 2000).

The $\dot{P}$ determinations reported in this section require confirmation with better data. This will soon be possible when OGLE III photometry is released. Combined OGLE II+III dataset will provide the timebase similar to that of MACHO, but much higher data quality should result in $\dot{P}$ errors about twice smaller.

Further accumulation of data will eventually allow determination of $\dot{P}$ for all three modes. This is quite essential for understanding the true nature of the period changes. If they are caused by stellar evolution, then all modes should yield roughly the same values of $\dot{P}/P$. No such equality is expected if period changes result from nonlinear mode interactions.

7. Conclusions and discussion

We show that pulsation periods measured in two LMC triple-mode Cepheids are consistent with values calculated for the first three overtones only if the envelope metal abundance is between $Z = 0.004$ and 0.007 . This is somewhat lower than $Z = 0.008$, typically adopted for the LMC. In one of the stars, fitting the periods requires the envelope hydrogen abundance $X \lesssim 0.72$, which is again somewhat below the commonly accepted LMC value of $X = 0.74$. The conclusions regarding envelope chemical composition rests mainly on the measured period ratios. For calculating the model periods, we rely on the linear pulsation theory. We believe, that nonlinear corrections to the periods are small, but we cannot estimate them. Therefore, it is important that the derived metallicity limits are verified through direct spectroscopic determination.

Effective temperatures implied by the models are consistent with observed $(V - I)$ colours of the two stars. Unfortunately, the accuracy of the latter is currently too low to yield useful constraints. Our models have luminosities confined to narrow ranges. The inferred distance moduli are $18^m34 - 18^m43$ for one star and $18^m49 - 18^m53$ for the other. The difference between the two objects can be in large part attributed to photometric errors. The derived distances to the triple-mode Cepheids are within the range currently considered for the LMC, but favour rather longer LMC distances in this range.

We find that parameters inferred from the pulsation periods of the two stars are consistent with evolutionary models undergoing the Hertzsprung gap crossing. The consistency does not require any overshooting from the convective core during the main sequence evolution. Seismic

models do not exclude overshooting, though, but rather impose an upper limit on its extent. The limit is particularly strong in case of LMC SC5-338399. The extent of overshooting can be further constrained once metal abundances or, alternatively, mean dereddened colours of the stars are precisely measured.

Both the periods and the luminosities of the triple-mode Cepheids cannot be reconciled with evolutionary models in the phase of central helium burning. There is nothing strange in our proposal that these stars are undergoing Hertzsprung gap crossing. Stars with masses of $\sim 3M_{\odot}$, which we find for our objects, spend about 10^5 years crossing the instability strip on their way to the red giant branch. This is comparable to time spent inside the strip by more massive stars ($M \sim 7M_{\odot}$) during the central helium burning phase. The pulsation periods of such stars are between 10 and 20 days. We observe many LMC Cepheids in this period range.

What may be a problem for the first crossing scenario is a possible conflict between predicted and measured rates of period change. The currently available data are insufficient for significant $\dot{P}$ determination. We note, however, that in both Cepheids the most probable $\dot{P}$ values are more than 3.8σ away from theoretical expectations. Since we are somewhat sceptical about $\dot{P}$ measurement errors, we do not yet consider this result to invalidate our models. But this is a warning sign.

If the future determinations confirm the discrepancy, our models will be in serious troubles. In our opinion, the problem would be more severe than the one encountered by Pietrukowicz (2003) for long period Cepheids. He has found that theoretically predicted rates of period change are faster than the measured ones, but the size of the effect depends on whose evolutionary tracks are used. According to our unpublished calculations, there is no significant conflict if one uses the tracks of Girardi et al. (2000).

The speed of evolution during the second and third crossings of the instability strip is affected by the element mixing in three distinct evolutionary phases. There are uncertainties regarding mixing efficiency in each of them. During the main sequence phase, we do not know the extent of the convective overshooting from the hydrogen burning core. During the red giant phase, we do not know the efficiency of the dredge up of the helium enriched material into the deep convective envelope. Finally, during the blue loop phase, there is a subtle problem with treatment of the boundary of the helium burning core. In the case of the first crossing of the instability strip, only the uncertainty of the main sequence overshooting affects the results, and this has been taken into account in our calculations. We believe that our prediction for the rates of period change is quite robust. Therefore, the possible discrepancy between calculated and observed $\dot{P}$ would pose a very serious challenge to our understanding of stellar evolution. Further observations of both triple-mode Cepheids are needed to settle this important issue.

Acknowledgements. This work has been supported in part by Polish KBN grant 5 P03D 030 20.

References

- Alibert, Y., Baraffe, I., Hauschildt, P. & Allard, F. 1999, *A&A*, 344, 551
- Allsman, R. A. & Axelrod, T. S. 2001, *astro-ph/0108444*
- Costa, J. E. S., Kepler, S. O. & Winget, D. E. 1999, *ApJ*, 522, 973
- Costa, J. E. S. & Kepler, S. O. 2000, *Baltic Astronomy*, 9, 451
- Dziembowski, W. 1977, *Acta Astron.*, 27, 95
- Fitch, W. S. & Szeidl, B. 1976, *ApJ*, 203, 616
- Fokin, A. B. 1994, *A&A*, 292, 133
- Girardi, L., Bressan, A., Bertelli, G. & Chiosi, C. 2000, *A&A*, 354, 892
- Iglesias, C. A. & Rogers, F. J. 1996, *ApJ*, 464, 943
- Kepler, S. O. 1993, *Baltic Astronomy*, 2, 515
- Kovács, G. 2000a, *A&A*, 360, L1
- Kovács, G. 2000b, *A&A*, 363, L1
- Kovács, G. & Buchler, J. R. 1988, *ApJ*, 324, 1026
- Kovács, G. & Buchler, J. R. 1994, *A&A*, 281, 749
- Kurucz, R. L. 1998, <http://kurucz.harvard.edu>
- Madore, B. F. & Freedman, W. L. 1991, *PASP*, 103, 933
- Moskalik, P. & Buchler, J. R. 1991, *ApJ*, 366, 300
- Moskalik, P. & Kołaczkowski, Z. 2005, in preparation
- Moskalik, P., Kołaczkowski, Z. & Mizerski, T. 2004, in *Variable Stars in the Local Group*, ed. D. W. Kurtz & K. Pollard, *ASP Conf. Ser.*, 310, 498
- Paczyński, B. 1970, *Acta Astron.*, 20, 47
- Pietrukowicz, P. 2003, *Acta Astron.*, 53, 63
- Popielski, B. L., Dziembowski, W. A., Cassisi, S., 2000, *Acta Astron.*, 50, 491
- Press, W. H., Teukolsky, S. A., Vetterling, W. T. & Flannery, B. P. 1986, *Numerical Recipes in FORTRAN* (2nd ed.; Cambridge: Cambridge Univ. Press)
- Rogers, F. J., Swenson, F. J. & Iglesias, C. A. 1996, *ApJ*, 456, 902
- Soszyński, I., Udalski, A., Szymański, M., et al. 2000, *Acta Astron.*, 50, 451
- Subramaniam A. 2003, *ApJL*, 598, L19
- Subramaniam A. 2004, *astro-ph/0409492*
- Udalski, A., Soszyński, I., Szymański, M., et al. 1999, *Acta Astron.*, 49, 223
- Żebruń, K., Soszyński, I., Woźniak, P. R. et al. 2001, *Acta Astron.*, 51, 317